

Editorial Comment

Dynamic Subaortic Obstruction: A Disease of the Mitral Valve Suitable for Surgical Repair?*

EHUD SCHWAMMENTHAL, MD,†
ROBERT A. LEVINE, MD, FACC*

Boston, Massachusetts and Tel Hashomer, Israel

Role of the mitral valve in subaortic obstruction. Because the left ventricle, in contrast to the right, has no separate inflow and outflow tract, it has to be converted from an inflow chamber in diastole to an outflow chamber in systole (1). This conversion is achieved by the anterior mitral leaflet switching like a relay from the septum toward the posterior wall, where it is tethered in systole by the papillary muscles. These muscles arise posteriorly and restrain the coapted leaflets below the outflow stream, which can proceed unimpeded; any disruption of this tethering mechanism, therefore, has the potential to allow the leaflets to move anteriorly and obstruct outflow. Such changes are well recognized to occur in hypertrophic cardiomyopathy, in which anterior displacement of the papillary muscles and mitral leaflet elongation diminish the effective tethering of the leaflets posteriorly and provide sufficient leaflet mobility to allow systolic anterior motion (SAM) of the leaflets, obstructing the outflow tract (2-11). Obstructive hypertrophic cardiomyopathy is therefore as much a disease of the mitral valve as it is of the myocardium. As a consequence, particularly in patients regarded as suboptimal candidates for septal myectomy, attempts have been directed to abolish outflow tract obstruction and mitral regurgitation by mitral valve surgery, the most definitive approach being excision of the native valve and replacement by a prosthesis (12). To avoid the limitations of prostheses, mitral valve plication has recently been introduced by McIntosh et al. (13) as an adjunctive procedure for patients judged at operation to be at increased risk for suboptimal hemodynamic results from septal myectomy alone. The decision to reduce anterior leaflet width by

plication is guided by intraoperative and preoperative echocardiography and direct observations of an enlarged and elongated anterior mitral leaflet persisting to prolapse into the outflow tract with septal contact despite apparently adequate muscular resection. This approach has yielded excellent hemodynamic and symptomatic results comparable to those of myectomy alone in patients without substantially enlarged mitral leaflets (13).

Anterior mitral leaflet extension. Whereas mitral valve plication reduces anterior leaflet size and thus follows the logic of pathoanatomic and echocardiographic studies of the mechanism of SAM, Kofflard et al. (14) in this issue of the Journal propose the exact opposite—horizontal anterior mitral leaflet extension—and with apparently no less impressive results. In a series of eight patients in whom the surgeon considered myectomy alone unlikely to yield optimal results, a glutaraldehyde-preserved autologous pericardial patch was inserted into the anterior leaflet, in a modification of a technique initially developed by Carpentier to treat mitral regurgitation in rheumatic valvular disease and bacterial endocarditis with perforation (15).

Why does it work? As suggested by the authors, the patch extends across the bending point of the buckling anterior leaflet and may stiffen it, preventing SAM. If this is the principal mechanism, simply sewing a pericardial patch onto the central portion of the leaflet would be expected to do the job without changing leaflet dimension. The basis for another explanation is the observation that SAM is typically greatest at the leaflet center (2,3,5,7), consistent with increased slack of the related chordae (e.g., created by inward or central displacement of the papillary muscle tips, as seen in hypertrophic obstructive cardiomyopathy). The authors therefore propose that inserting the central patch may displace the chordae inserting into the central leaflet laterally, thus stretching them and increasing leaflet tethering. Further, taking the anterior leaflet margins, which have been incised along a straight line, and deforming them to follow the longer elliptical outline of the patch should make the leaflet slightly shorter because leaflet area and mass remain the same; this change would therefore reduce the slack needed to produce SAM.

An alternative hypothesis would be the additive beneficial impact of increased leaflet area once the streamlines of flow have been normalized by septal myectomy. In the presence of a hypertrophied basal anterior septum bulging into the cavity, blood flow cannot take the direct route from the apex to the outflow tract, but must circumvent the septum (Fig. 1). The streamlines of flow are therefore curved and may hit the anterior leaflet on its posterior surface, pushing it toward the septum. Once normalization (straightening) of the intraventricular streamlines has been achieved by subvalvular myectomy, flow is directed onto the anterior surface of the mitral leaflets, as in normal hearts; in this case, increasing leaflet area by horizontal extension (which probably does not increase leaflet slack) will be beneficial in preventing SAM and mitral regurgitation because the anterior leaflet is then pressed

*Editorials published in *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of JACC or the American College of Cardiology.

From the *Cardiac Ultrasound Laboratory, Massachusetts General Hospital and Department of Medicine, Harvard Medical School, Boston, Massachusetts; and †Cardiac Ultrasound Laboratory, Chaim Sheba Medical Center, Tel Aviv University, Tel Hashomer, Israel.

This work was supported in part by Grants HL38176 and HL53702 from the National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, Maryland and by a grant from the Israel Heart Association. Dr. Levine is an Established Investigator of the American Heart Association, Dallas, Texas, with funds supplied in part by its Massachusetts Affiliate, Needham, Massachusetts.

Address for correspondence: Dr. Ehud Schwammenthal, Cardiac Ultrasound Laboratory, Chaim Sheba Medical Center, 52621 Tel Hashomer, Israel.




Figure 1. Top. In addition to the force generated by the papillary muscles (directed apically and posteriorly, as shown by the arrows below the mitral valve), the flow that is directed onto the anterior surface of the mitral leaflet contributes to keeping the mitral valve posteriorly with a force that equals left ventricular pressure times the area of the leaflet exposed to that pressure. **Bottom.** Because of a hypertrophied basal anterior septum bulging into the cavity, blood flow cannot take the direct path from the apex to outflow tract, but necessarily must circumvent the septum. The streamlines of flow are therefore curved and may hit the anterior leaflet on its posterior surface, dragging it toward the septum; leaflet mobility to follow the forces of the flow field is facilitated by abnormal mobility of the mitral valve due to leaflet elongation and anterior displacement of the papillary muscles. AML = anterior mitral leaflet; Ao = aorta; LA = left atrium; LV = left ventricle; HOCM = obstructive hypertrophic cardiomyopathy; IVS = interventricular septum; PML = posterior mitral leaflet; Po = posterior wall of the left ventricle.

posteriorly by a force that is proportional to the leaflet area exposed to flow. In fact, a flow stream directly impacting on the anterior surface of a tethered mitral leaflet in vitro actually maintains its posterior position and effectively opposes the lift or Venturi force created by a higher velocity outflow stream not impacting on the leaflet surface (16). The practical implication of this mechanism is that horizontal leaflet extension would be helpful after septal myectomy (in a synergistic manner) but would not necessarily reduce SAM without it, whereas mitral leaflet plication has the potential to reduce or even abolish SAM independently; leaflet stiffening caused by the patch, of course, would reduce SAM independent of myectomy as well.

Another observation of mechanistic interest is that mitral regurgitation, which was more severe before operation in the group undergoing leaflet extension, was abolished more effectively by providing greater leaflet area; this is consistent with

previous observations, both in vitro and in patients, that increased leaflet contact area reduces mitral regurgitation for a given degree of SAM (11,17,18).

Does it work? Proving the value of an additional mitral valve intervention after myectomy would require either demonstrating more effective results with than without mitral valve surgery (either in the same patients or in matched groups, not chosen in the present study) or demonstrating good hemodynamic results with additional mitral valve surgery in patients known to do poorly with myectomy alone, as the authors set out to do. It is not clear, however, that such proof is available on the basis of the selection process in this study, which related to the impression regarding expected hemodynamic outcome. Obviously, the severity of mitral regurgitation favored mitral valve extension. Other conditions generally favoring mitral valve surgery, such as limited septal hypertrophy, although mentioned in the methods and discussion sections, did not appear to play an actual role: There was no difference in the range of septal thickness in the group with and without additional mitral leaflet extension (20 to 35 mm vs. 20 to 33 mm). Moreover, in the group undergoing additional mitral leaflet extension, myectomy was significantly more effective in reducing septal thickness (19 ± 4 mm vs. 22 ± 4 mm, $p < 0.05$, Table 2 [14]). In view of the limited number of patients, as the authors note, it cannot be excluded that at least some of the beneficial effect attributed to the additional mitral leaflet extension was actually caused by a more effective myectomy in the leaflet extension group. No information is provided as to whether there was a difference in mean mitral leaflet area between the two groups, although it is mentioned that all patients in both groups had clearly enlarged leaflets, and whether differences in leaflet length and area could potentially be valuable for selecting the appropriate procedure. Finally, it is not clear why postoperative functional class was better in the leaflet extension group despite a comparable hemodynamic result. The differences in functional class cannot all be related to decreased residual mitral regurgitation (Table 1 [14]). It is even conceivable that the effect of medication (such as antiarrhythmic agents or a combination of beta-adrenergic blocking agents and calcium antagonists, which has been shown to increase pulmonary artery pressure significantly without any beneficial effect compared with each drug alone [19]) may have contributed to the decreased functional capacity of the patients undergoing myectomy alone, who were taking more medications postoperatively (Table 2 [14]). A placebo effect in patients who know they have undergone an "improved" procedure also cannot be excluded without objective exercise testing, which can safely be performed in these patients (20,21), particularly after relief of outflow tract obstruction.

Is it necessary? Subvalvular myectomy has been shown to be an effective treatment for symptomatic patients with obstructive hypertrophic cardiomyopathy refractory to medical treatment that can be performed with low mortality and morbidity in experienced hands, yielding excellent symptomatic and hemodynamic long-term results, with an experience of >25 years (22-26). A subset of patients, however, may be

difficult to manage without additional mitral valve surgery, even in experienced hands (13,14), and could therefore benefit from mitral leaflet plication or extension. Definite proof of the superiority of combined myectomy and mitral valve repair would require a staged approach, with assessment of hemodynamic variables by intraoperative echocardiography after myectomy alone (off bypass) and then after the additional mitral valve repair. (Such multiple bypass times are routinely used in some centers [27] for optimizing myectomy by intraoperative echocardiography and thus do not involve the ethical problems posed by a randomized study in this context.) Quantification of mitral leaflet geometry by intraoperative echocardiography could potentially help us to develop clearly defined indications for mitral valve repair as well as guide the procedure. Nevertheless, the study by Kofflard et al. (14) is consistent with growing evidence that structural abnormalities of the mitral apparatus can play a primary role in causing SAM and dynamic subaortic obstruction (2-11,28,29). In hypertrophic cardiomyopathy, SAM can be viewed as the result of an abnormal mitral valve apparatus exposed to an abnormal flow field; leaflet extension after myectomy addresses both of these factors. Systolic anterior motion after mitral valve repair for myxomatous prolapse can also be related to slack leaflet portions interposed into the outflow stream because the posterior leaflet is elongated and shifts the coaptation point anteriorly (30-33). These observations are consistent with lessons learned from experimental and computational models studying the interaction of the mitral valve and outflow stream (5-7,16,34-38). This increased understanding has direct practical implications for surgical management, and it is our expectation that detailed cardiac ultrasound imaging of the underlying anatomy (39) will allow new techniques to be applied most effectively to the individual patient to provide the most normal function and the greatest benefit.

We thank Karen Reynolds, MPH, for expert assistance.

References

- Silverman ME, Hurst JW. The mitral complex: interaction of the anatomy, physiology, and pathology of the mitral annulus, mitral valve leaflets, chordae tendineae, and papillary muscles. *Am Heart J* 1968;76:399-418.
- Jiang L, Levine RA, King ME, Weyman AE. An integrated mechanism for systolic anterior motion of the mitral valve in hypertrophic cardiomyopathy based on echocardiographic observations. *Am Heart J* 1987;113:633-44.
- Maron BJ, Harding AM, Spirito P, Roberts WC, Waller BF. Systolic anterior motion of the posterior mitral leaflet: a previously unrecognized cause of dynamic subaortic obstruction in patients with hypertrophic cardiomyopathy. *Circulation* 1983;68:282-93.
- Shah P, Taylor RD, Wong M. Abnormal mitral valve coaptation in hypertrophic obstructive cardiomyopathy: proposed role in systolic anterior motion of mitral valve. *Am J Cardiol* 1981;48:258-62.
- Cape EG, Simons D, Jimoh A, Weyman AE, Yoganathan AP, Levine RA. Chordal geometry determines the shape and extent of systolic anterior mitral motion: in vitro studies. *J Am Coll Cardiol* 1989;13:1438-48.
- Sherrid MV, Chu CK, Delia E, Mogtader A, Dwyer EM. An echocardiographic study of the fluid mechanics of obstruction in hypertrophic cardiomyopathy. *J Am Coll Cardiol* 1993;22:816-25.
- Levine RA, Vlahakes GJ, Lefebvre X, et al. Papillary muscle displacement causes systolic anterior motion of the mitral valve. Experimental validation and insights into the mechanism of subaortic obstruction. *Circulation* 1995;91:1189-95.
- Klues HG, Maron BH, Dollar AL, Roberts WC. Diversity of structural mitral valve alterations in hypertrophic cardiomyopathy. *Circulation* 1992;85:1651-60.
- Grigg LE, Wigle ED, Williams WG, Daniel LB, Rakowski H. Transesophageal Doppler echocardiography in obstructive hypertrophic cardiomyopathy: clarification of pathophysiology and importance in intraoperative decision making. *J Am Coll Cardiol* 1992;20:42-52.
- Klues HG, Dollar AL, Prosser MA, et al. Echocardiographic assessment of mitral valve size in obstructive hypertrophic cardiomyopathy: anatomical validation from mitral valve specimens. *Circulation* 1993;88:548-55.
- Klues HG, Roberts WC, Maron BJ. Morphological determinants of echocardiographic patterns of mitral valve systolic anterior motion in hypertrophic obstructive cardiomyopathy. *Circulation* 1993;87:1570-9.
- Krajcer Z, Leachman RD, Cooley DA, Coronado R. Septal myotomy-myomectomy versus mitral valve replacement in hypertrophic cardiomyopathy: ten-year follow-up in 185 patients. *Circulation* 1989;80 Suppl I:1-57-64.
- McIntosh CL, Maron BJ, Cannon RO, Klues HG. Initial results of combined anterior mitral leaflet plication and ventricular septal myotomy-myomectomy for relief of left ventricular outflow tract obstruction in patients with hypertrophic cardiomyopathy. *Circulation* 1992;86 Suppl II:1-160-7.
- Kofflard MJ, van Herwerden LA, Waldstein DJ, et al. Initial results of combined anterior mitral leaflet extension and myectomy in patients with obstructive hypertrophic cardiomyopathy. *J Am Coll Cardiol* 1996;28:197-22.
- Chauvaud S, Jebara V, Chachques JC, et al. Valve extension with glutaraldehyde-preserved autologous pericardium. *J Thorac Cardiovasc Surg* 1991;102:171-8.
- Simpson MS, Lefebvre XP, He S, Yoganathan AP, Levine RA. Outflow impingement on the mitral valve counteracts the Venturi effect and prevents systolic anterior motion: flow studies [abstract]. *Circulation* 1991;84 Suppl II:II-146.
- Schwammenthal E, He S, Hopmeyer J, et al. Mechanism of mitral regurgitation in hypertrophic cardiomyopathy: competition between systolic anterior motion of the posterior leaflet and its restraint [abstract]. *J Am Coll Cardiol* 1993;21:353A.
- Schwammenthal E, Nakatani S, Lever HM, Weyman AE, Levine RA. Intraoperative transesophageal studies of the mechanism of mitral regurgitation in hypertrophic cardiomyopathy [abstract]. *Circulation* 1994;90(Pt 2):I-170.
- Hopf R, Kaltenbach M. Ten-year results and survival of patients with hypertrophic cardiomyopathy treated with calcium antagonists. *Zeitschrift für Kardiologie* 1987;76 Suppl 3:137-44.
- Lösse B, Kuhn H, Loogen F, Schulte D. Exercise performance in hypertrophic cardiomyopathy. *Eur Heart J* 1983;4 Suppl F:197-208.
- Schwammenthal E, Schwartzkopff B, Block M, et al. Doppler echocardiographic assessment of the pressure gradient during bicycle ergometry in hypertrophic cardiomyopathy. *Am J Cardiol* 1992;69:1623-8.
- Morrow AG, Reitz BA, Epstein SE, et al. Operative treatment in hypertrophic subaortic stenosis: techniques and the results of pre- and postoperative assessments in 83 patients. *Circulation* 1975;88-102.
- Maron BJ, Epstein SE, Morrow AG. Symptomatic status and prognosis of patients after operation for hypertrophic obstructive cardiomyopathy: efficacy of ventricular septal myotomy and myectomy. *Eur Heart J* 1983;4 Suppl F:175-85.
- Williams WG, Wigle ED, Rakowski H, et al. Results of surgery for hypertrophic obstructive cardiomyopathy. *Circulation* 1987;76 Suppl V-V:104-8.
- Bircks W, Schulte HD. Surgical treatment of hypertrophic obstructive cardiomyopathy with special reference to complications and to atypical hypertrophic cardiomyopathy. *Eur Heart J* 1983;4 Suppl F:187-90.
- Seiler C, Hess OM, Schoenbeck M, et al. Long-term follow-up of medical versus surgical therapy for hypertrophic cardiomyopathy: a retrospective study. *J Am Coll Cardiol* 1991;17:634-42.
- Marwick TH, Stewart WJ, Lever HM, et al. Benefits of intraoperative echocardiography in the surgical management of hypertrophic cardiomyopathy. *J Am Coll Cardiol* 1992;20:1066-72.

28. Moro E, Ten Cate JH, Hugenholtz PG, Roelandt J. Genesis of systolic anterior motion of the mitral valve in hypertrophic cardiomyopathy: an anatomical or dynamic event? *Eur Heart J* 1987;8:1312-21.
29. Nakatani S, Schwammenthal E, Lever HM, Levine RA, Lytle BW, Thomas JD. New insights into the reduction of mitral valve systolic anterior motion after ventricular septal myectomy in hypertrophic obstructive cardiomyopathy. *Am Heart J* 1996;131:294-300.
30. Galler M, Kronzon I, Slater J, et al. Long-term follow-up after mitral valve reconstruction: incidence of postoperative left ventricular outflow obstruction. *Circulation* 1986;74 Suppl 1:I-99-103.
31. Mihaileanu S, Marino JP, Chauvaud S, et al. Left ventricular outflow obstruction after mitral valve repair (Carpentier's technique): proposed mechanisms of disease. *Circulation* 1988;78 Suppl 1:I-78-84.
32. Jebara VA, Mihaileanu SA, Acar C, et al. Left ventricular outflow obstruction after mitral valve repair: results of the sliding leaflet technique. *Circulation* 1993;88(Pt 2):30-4.
33. Lee KS, Stewart WJ, Lever HM, Underwood PL, Cosgrove DM. Mechanism of outflow tract obstruction causing failed mitral valve repair: anterior displacement of leaflet coaptation. *Circulation* 1993;88(Pt 2):24-9.
34. Lefebvre XP, Yoganathan AP, Levine RA. Insights from in-vitro flow visualization into the mechanism of systolic anterior motion of the mitral valve in hypertrophic cardiomyopathy under steady flow conditions. *J Biomech Eng* 1992;114:406-13.
35. Lefebvre XP, He S, Levine RA, Yoganathan AP. Systolic anterior motion of the mitral valve in hypertrophic cardiomyopathy: an in vitro pulsatile flow study. *J Heart Valve Dis* 1995;4:422-38.
36. Yoganathan AP, Lemmon JD, Kim YH, Levine RA, Vesier CC. A three-dimensional computational investigation of intraventricular fluid dynamics: examination into the initiation of systolic anterior motion of the mitral valve leaflets. *J Biomech Eng* 1995;117:94-102.
37. He S, Hopmeyer J, Lefebvre X, Yoganathan AP, Levine RA. Importance of leaflet elongation in causing systolic anterior motion of the mitral valve [abstract]. *Circulation* 1992;86 Suppl 1:I-850.
38. Levine RA, Lefebvre X, Cuerrero JL, et al. Unifying concepts of mitral valve function and disease: SAM, prolapse and ischemic mitral regurgitation. *J Cardiol* 1994;24 Suppl 38:157-69.
39. Maslow AE, Levine RA. Echocardiographic predictors of left ventricular outflow tract obstruction and systolic anterior motion of the mitral valve after mitral valve reconstruction for myxomatous valve disease [abstract]. *J Am Coll Cardiol* 1996;27 Suppl A:232A.